

AD-A055 563

CENTER FOR NAVAL ANALYSES ARLINGTON VA
CORRESPONDENCE RULES AND PATH INTEGRALS. (U)

F/G 12/1

UNCLASSIFIED

JUN 78 M M MIZRAHI
CNA-PP-222

NL

1 of 1
AD
A055 563

END



END
DATE
FILMED
8-78
DDC

AD A 055563

AD No. _____
FILE COPY

55 000222.00

2

6 CORRESPONDENCE RULES AND PATH INTEGRALS,

9 Professional paper,

10 Maurice M. Mizrahi

Professional Paper No. 222

11 June 1978

12 37 p.

14 CNA-PP-222

The ideas expressed in this paper are those of the author. The paper does not necessarily represent the views of the Center for Naval Analyses.

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

DDC
RECEIVED
JUN 26 1978
B

CENTER FOR NAVAL ANALYSES

1401 Wilson Boulevard
Arlington, Virginia 22209

78 06 16 030

077 270

JOB

CORRESPONDENCE RULES AND PATH INTEGRALS

Maurice M. Mizrahi

Center for Naval Analyses of
the University of Rochester
1401 Wilson Boulevard
Arlington, Va. 22209, USA

78 06 16 030

ABSTRACT

A path-integral representation is constructed for propagators corresponding to quantum Hamiltonian operators obtained from classical Hamiltonians by an arbitrary rule of correspondence. Each rule yields a unique way of defining the path integral in the context of a formalism which does not require a limiting process. This formalism is more reliable than the usual lattice definition in that all the expressions it entails are well-defined for computational purposes and it allows the explicit evaluation of large classes of path integrals. Direct substitution in the Schrödinger equation shows that there are no restrictions on the Hamiltonian operator. Examples are given.

ACCESSION		
NTIS	WALL	✓
DDO	ALL	□
UNCLASSIFIED		□
PER FORM 50		
BY		
DISTRIBUTION/AVAILABILITY CODES		
Or: 100/100 SPECIAL		
A		

I. INTRODUCTION

The purpose of this paper is to propose a solution to the following problem: Given an arbitrary classical Hamiltonian $H_c(p,q,t)$ and an arbitrary rule of correspondence which enables one to derive a quantum Hamiltonian operator $\underline{H}(P,Q,t)$ (Hermitian or not) from H_c , find a path-integral representation for the propagator $K \equiv \langle q_b, t_b | q_a, t_a \rangle$ corresponding to \underline{H} which (1) takes proper account of the correspondence rule and (2) does not involve a limiting process (i.e. a "skeletonization of the path" or "time-slicing" technique) in its definition. The latter requirement purports to avoid the many ambiguities inherent in this process and to enable one to actually compute path integrals, rather than simply exhibit formal expressions.

In 1975 we showed¹, by time-slicing and Weyl transform techniques, that a formal path-integral expression in phase space can be written for the propagator, where the Weyl transform of the Hamiltonian operator takes the place of the classical Hamiltonian in the action functional. Subsequent work by Cohen² and Dowker³ showed that formal path integrals can be obtained to accommodate any rule of correspondence. In all the foregoing papers, however, work stopped when a "formal" path integral was obtained, leaving open the problems of evaluation of the path integrals, substitution of the path-integral expression in the Schrödinger equation for verification, and justification of some possibly ambiguous limits inherent in the time-slicing approach. This paper will address the above problems by proposing an alternative approach, and supersedes Ref. 1.

The starting points are the general framework for path integration in phase space without limiting procedure introduced in Ref. 4 and Cohen's

mappings between correspondence rules and ordinary functions⁵. It will be shown that an infinite series representing the propagator (where each term is a path integral) satisfies the Schrödinger equation with arbitrary \underline{H} and the boundary condition, provided a consistent well-defined algorithm is used to properly take the correspondence rule into account.

Only path integrals representations with respect to the free-particle measure will be considered here. The more general measures introduced in Ref. 4, which allow a semiclassical expansion of the propagator, will be treated elsewhere.⁶ We work in one dimension to simplify the discussion, although the results can be readily generalized to n dimensions. All integrals are over \mathbb{R}^s for suitable s , unless otherwise specified.

II. THE PATH-INTEGRAL FORMALISM

The formalism for constructing phase-space path integrals without resorting to a limiting procedure with ambiguous epsilonics was introduced in Ref. 4 and only a brief description will be given here⁷. It consists of defining what plays the role of a measure in phase space by its Fourier transform, which is a simple closed-form expression. For example, the normalized free-particle measure $w(p,q)$ in phase space \mathcal{P} , corresponding to

$$dw(p,q) \sim \frac{1}{K_0} \left[\frac{dp dq}{2\pi\hbar} \right] \exp \left\{ \frac{i}{\hbar} \int_{t_a}^{t_b} \left[p(t) \dot{q}(t) - \frac{p^2(t)}{2m} \right] dt \right\}, \quad (1)$$

can be shown⁴ to have, as its Fourier transform:

$$F_w(\mu, \nu) = \exp \left\{ -i \langle \mu, \bar{q} \rangle - i \langle \nu, \bar{p} \rangle - \frac{i\hbar}{2} \iint_T G_{ab}(t, t') d\mu(t) d\mu(t') \right. \\ \left. - i\hbar \iint_T \bar{G}(t, t') d\mu(t) d\nu(t') - \frac{i\hbar}{2} \iint_T G_p(t, t') d\nu(t) d\nu(t') \right\}, \quad (2)$$

where:

$$(1) \mathcal{P} \equiv \left\{ [p(t), q(t)] \text{ on } T \equiv [t_a, t_b] \mid q(t_a) = q_a, q(t_b) = q_b, p(t) \text{ unrestricted} \right\}, \quad (3)$$

$$(2) K_0 = \left(\frac{m}{2\pi i \hbar T} \right)^{1/2} \exp \left[\frac{im(q_b - q_a)^2}{2\hbar T} \right] \quad (4)$$

is the free-particle propagator ($T \equiv t_b - t_a$),

(3) $[\bar{q}(t), \bar{p}(t)]$ is the average free-particle path:

$$\bar{q}(t) = [q_b(t - t_a) + q_a(t_b - t)] / T \quad (5)$$

$$\bar{p}(t) = m(q_b - q_a) / T \quad (6)$$

(4) The G functions are the free-particle covariances:

$$G_{ab}(t, t') = [(t' - t_a)(t_b - t) \gamma(t - t') + (t - t_a)(t_b - t') \gamma(t' - t)] / mT \quad (7)$$

$$\bar{G}(t, t') = [(t_b - t) \gamma(t - t') - (t - t_a) \gamma(t' - t)] / T \quad (8)$$

$$G_p(t, t') = -m/T \quad (9)$$

$[\gamma(x) \text{ being } 1 \text{ when } x > 0 \text{ and } 0 \text{ otherwise}],$

(5) μ and ν are elements of \mathcal{M} , the space of bounded measures on the time interval T ; $\langle \mu, q \rangle \equiv \int_T q(t) d\mu(t)$ if μ is induced by a function, $\langle \delta_t, q \rangle \equiv q(t)$ if μ is δ_t , the "delta-function" measure at t .

The most general Gaussian measure, which absorbs all the quadratic terms [i.e. not only $p^2/2m$ but $g(t)p^2/2m + f(t)q^2/2 + k(t)pq$] in either the Hamiltonian or the action functional expanded about the classical path, was constructed in Ref. 4, but the free-particle measure will be sufficient for our purpose here. w is a true phase-space measure: it does not entail performing separate, successive path integrals in configuration space and momentum space.

This definition enables one to carry out path integrals of cylindrical functionals, i.e. functionals which depend on only a finite number of terms of the form $\langle \mu, q \rangle$ or $\langle \nu, p \rangle$, by converting them into finite-dimensional ^{ordinary} integrals. The result is the following fundamental integral [in a form slightly different from that given in Ref. 4, Eq. (97)]:

$$\begin{aligned} & \int_{\mathcal{P}} F(\langle \mu_1, q \rangle, \dots, \langle \mu_n, q \rangle, \langle \nu_1, p \rangle, \dots, \langle \nu_m, p \rangle) dw(p, q) \\ &= \int_{\mathbb{R}^{n+m}} \frac{F(y) dy}{[(2\pi i \hbar)^{m+n} \det A]^{1/2}} \exp \left\{ \frac{i}{2\hbar} \sum_{i,j=1}^{n+m} (A^{-1})_{ij} (y_i - a_i)(y_j - a_j) \right\} \end{aligned} \quad (10)$$

where

$$y \equiv (u_1, \dots, u_n, v_1, \dots, v_m) \quad (11)$$

$$dy \equiv du_1 \dots du_n dv_1 \dots dv_m \quad (12)$$

$$a \equiv (\langle \mu_1, \bar{q} \rangle, \dots, \langle \mu_n, \bar{q} \rangle, \langle \nu_1, \bar{p} \rangle, \dots, \langle \nu_m, \bar{p} \rangle) \quad (13)$$

$$A \equiv \begin{pmatrix} W & C \\ \tilde{C} & V \end{pmatrix} \quad [(n+m) \times (n+m)] \quad (14)$$

$$W_{ij} \equiv \int_T \int_T G_{ab}(t, t') d\mu_i(t) d\mu_j(t') \quad (n \times n) \quad (15)$$

$$C_{ij} \equiv \tilde{C}_{ji} \equiv \int_T \int_T \bar{G}(t, t') d\mu_i(t) dv_j(t') \quad (n \times m) \quad (16)$$

$$V_{ij} \equiv \int_T \int_T G_p(t, t') dv_i(t) dv_j(t') \quad (m \times m) \quad (17)$$

In the above formula, w is not restricted to being the free-particle measure, but can be the most general quadratic measure mentioned earlier. Of particular interest are an expression for the Fourier transform of w :

$$\int_{\mathcal{P}} \left\{ \exp[-i\langle \mu, q \rangle - i\langle \nu, p \rangle] \right\} dw(p, q) = \bar{F}w(\mu, \nu) \quad (18)$$

and the special case where $F(x_1, \dots, x_k) = x_1 \dots x_k$, which yields the generalized moments formula in phase space [an extension of the one given in Ref. 8, Eq. (65), for configuration space] :

$$\begin{aligned} & \int_{\mathcal{P}} \langle \mu_1, q \rangle \dots \langle \mu_n, q \rangle \langle \nu_1, p \rangle \dots \langle \nu_m, p \rangle dw(p, q) \\ &= i^{n+m} \mathcal{H}_{n+m} \left(-\frac{ic_1}{2}, \dots, -\frac{ic_{n+m}}{2} \right) \end{aligned} \quad (19)$$

where

$$c_k \equiv \begin{cases} \langle \mu_k, \bar{q} \rangle & \text{for } k = 1, \dots, n \\ \langle \nu_k, \bar{p} \rangle & \text{for } k = n+1, \dots, n+m \end{cases} \quad (20)$$

and \mathcal{H}_{n+m} is the generalized Hermite polynomial of order $n+m$ and matrix $i\hbar A/2$, defined in Refs. 8 and 9¹⁰. The examples shown below, for the case where all μ s and ν s are δ functions, reveal the general pattern:

$$\int_{\mathcal{P}} q(t) dw(p, q) = \bar{q}(t) \quad (21)$$

$$\int_{\mathcal{P}} q(t) p(t') dw(p, q) = \bar{q}(t) \bar{p}(t') + i\hbar \bar{G}(t, t') \quad (22)$$

$$\int_{\mathcal{P}} q(t) q(t') dw(p, q) = \bar{q}(t) \bar{q}(t') + i\hbar G_{ab}(t, t') \quad (23)$$

$$\int_{\mathcal{P}} p(t) p(t') dw(p, q) = \bar{p}(t) \bar{p}(t') + i\hbar G_p(t, t') \quad (24)$$

$$\begin{aligned} \int_{\mathcal{P}} q(t_1) q(t_2) p(t_3) dw(p, q) &= \bar{q}(t_1) \bar{q}(t_2) \bar{p}(t_3) \\ &+ i\hbar \bar{G}(t_2, t_3) \bar{q}(t_1) + i\hbar \bar{G}(t_1, t_3) \bar{q}(t_2) \\ &+ i\hbar G_{ab}(t_1, t_2) \bar{p}(t_3) \end{aligned} \quad (25)$$

$$\begin{aligned}
\int_{\mathcal{P}} q(t_1) q(t_2) p(t_3) p(t_4) d\omega(p, q) &= \bar{q}(t_1) \bar{q}(t_2) \bar{p}(t_3) \bar{p}(t_4) \\
&+ i\hbar \bar{G}(t_1, t_4) \bar{q}(t_2) \bar{p}(t_3) + i\hbar \bar{G}(t_2, t_4) \bar{q}(t_1) \bar{p}(t_3) \\
&+ i\hbar G_p(t_3, t_4) \bar{q}(t_1) \bar{q}(t_2) + i\hbar G_{ab}(t_1, t_2) \bar{p}(t_3) \bar{p}(t_4) \\
&+ i\hbar \bar{G}(t_2, t_3) \bar{q}(t_1) \bar{p}(t_4) + i\hbar \bar{G}(t_1, t_3) \bar{q}(t_2) \bar{p}(t_4) \\
&+ (i\hbar)^2 G_{ab}(t_1, t_2) G_p(t_3, t_4) + (i\hbar)^2 \bar{G}(t_2, t_3) \bar{G}(t_1, t_4) \\
&+ (i\hbar)^2 \bar{G}(t_1, t_3) \bar{G}(t_2, t_4)
\end{aligned} \tag{26}$$

It will be noted that the pq correlation function, $\bar{G}(t, t')$, is discontinuous across the diagonal $t = t'$, and its jump there is of magnitude 1:

$$\left[\lim_{(t-t') \rightarrow 0^+} - \lim_{(t'-t) \rightarrow 0^+} \right] \bar{G}(t, t') = 1 \tag{27}$$

[see, e.g., (8) for the free-particle case]. Thus, $\mathcal{F}\omega(\delta_t, \delta_t)$ is not defined. This indefiniteness, which occurs in a natural manner as one builds the measure⁴, stems from the non-commutativity of \underline{p} and \underline{q} , and will give us the flexibility we need to take various correspondence rules into account.

III. THE CORRESPONDENCE RULES

We will consider the most general quantum Hamiltonian operator $\hat{H}(\hat{P}, \hat{Q}, t)$ which can be derived from a classical Hamiltonian $H_c(p, q, t)$. The form most convenient for our purposes is that given by Cohen⁵:

$$\begin{aligned} \hat{H} = (2\pi\hbar)^{-2} \int dp dq du dv F(u, v, \hbar) H_c(p, q, t) \\ \times \exp \left\{ (i/\hbar) \left[(q - \hat{Q})u + (p - \hat{P})v \right] \right\}, \end{aligned} \quad (28)$$

where $F(u, v, \hbar)$ is the transformation function¹¹. Each F uniquely determines the correspondence rule. For instance, $F = 1$, $F = \cos(uv/2\hbar)$ and $F = (uv/2\hbar)^{-1} \sin(uv/2\hbar)$ give the Weyl, symmetrized, and Born-Jordan ordering schemes, respectively. For real H_c , \hat{H} is Hermitian iff $F^*(-u, -v, \hbar) = F(u, v, \hbar)$, a condition we will not need. Also, we must have $F(0, v, \hbar) = F(u, 0, \hbar) = 1$ to insure that $f(\hat{P})$ and $g(\hat{Q})$ correspond to $f(p)$ and $g(q)$. The transform can be inverted to yield

$$\begin{aligned} H_c(p, q, t) = (2\pi\hbar)^{-1} \int dq' dp' dq'' \exp \left\{ (i/\hbar) \left[q'p + p'(q - q') \right] \right\} \\ \times F^{-1}(q', p', \hbar) \left\langle q'' - \frac{1}{2}q' \right| \hat{H} \left| q'' + \frac{1}{2}q' \right\rangle, \end{aligned} \quad (29)$$

and finally, the transformation function F can be inferred from the knowledge of the Hamiltonian H_c and its transform \hat{H} :¹²

$$F(u, v, \hbar) \int dp dq H_c(p, q, t) \exp [i(qu + pv)/\hbar] \\ = (2\pi\hbar) \text{tr} \left(e^{i(\underline{Q}u + \underline{P}v)/\hbar} \underline{H} \right). \quad (30)$$

Note that the F which relates a given \underline{H} with a given H_c is usually not unique. For example, for $\underline{H} = f(\underline{Q})$ and $H_c = f(q)$, (30) gives:

$$F(u, v, \hbar) \delta(v) \int dq f(q) e^{iqu/\hbar} = \int dq f(q) e^{iqu/\hbar}. \quad (31)$$

Thus, any F such that $F(u, 0, \hbar) = 1$ will do. Similarly, one finds that pq is mapped into $(\underline{P}\underline{Q} + \underline{Q}\underline{P})/2$ by any $F = F(uv/\hbar)$ such that $F(0) = 1$ and $F'(0) = 0$.

It will be more convenient in certain cases to put \underline{H} in (28) in its normal-ordered form (\underline{Q} before \underline{P}). For this, one uses the Baker-Campbell-Hausdorff formula, $e^{\underline{A}+\underline{B}} = e^{\underline{A}}e^{\underline{B}}e^{-[\underline{A}, \underline{B}]/2}$, valid for all \underline{A} and \underline{B} which commute with $[\underline{A}, \underline{B}]$. It gives:

$$e^{-i(\underline{Q}u + \underline{P}v)/\hbar} = e^{-i\underline{Q}u/\hbar} e^{-i\underline{P}v/\hbar} e^{iuv/2\hbar}. \quad (32)$$

Since \underline{H} will be applied to functions of the endpoint q_b , \underline{Q} is represented by q_b and \underline{P} by $-\hbar \partial / \partial q_b$. Thus, the most general operator \underline{H} derived from H_c is, in its normal-ordered form,

$$\underline{H} = (2\pi\hbar)^{-2} \int dp dq du dv F(u, v, \hbar) H_c(p, q, t) \\ \times \exp \left[\frac{i u}{\hbar} \left(q - q_b + \frac{v}{2} \right) \right] \exp \left(\frac{i p v}{\hbar} \right) \exp \left(-v \frac{\partial}{\partial q_b} \right). \quad (33)$$

Consequently, when \underline{H} is applied to $f(q_b)$, the result is the right-hand side of (33) with $\exp(-v \partial / \partial q_b)$ replaced by $f(q_b - v)$.

IV. THE PROPAGATOR BY PATH INTEGRALS, FOR ARBITRARY HAMILTONIANS

With the two foregoing tools in hand, we can proceed to write a path-integral representation for the propagator corresponding to an arbitrary Hamiltonian operator. First, we write the latter in the form:

$$\underline{H} = \frac{\underline{P}^2}{2m} + k \underline{H}_1(\underline{P}, \underline{Q}, t). \quad (34)$$

This is done to split off the free-particle part so we can use the free-particle measure $w(p, q)$. Although the propagator will be expressed as a power-series in k , our aim is not a perturbation expansion. Rather, it is the manner in which the ordering of the factors in \underline{H}_1 is taken into account in evaluating the path integrals, so that the propagator K thereby obtained satisfies the Schrödinger equation associated with \underline{H} to all orders in k . Thus, k should be regarded as simply a "bookkeeping" parameter. Second, we define a function $\underline{H}_{no}(p, q, t)$, obtained by normal-ordering the operator \underline{H}_1 and then replacing \underline{P} by p and \underline{Q} by q . For example (since $\underline{Q}\underline{P} - \underline{P}\underline{Q} = i\hbar$):

$$\underline{H}_c = \underline{P}^2/2m + k p q^2 \quad (35)$$

$$\underline{H}_1 = \underline{P}^2/2m + k \underline{Q}\underline{P}\underline{Q} = \underline{P}^2/2m + k(\underline{Q}^2\underline{P} - i\hbar\underline{Q}) \quad (36)$$

$$\underline{H}_{no} = q^2 p - i\hbar q. \quad (37)$$

Our main result is expressed in the following theorem.

Theorem

The propagator $K \equiv \langle q_b, t_b | q_a, t_a \rangle$, or probability amplitude that a particle at position q_a at time t_a will be at position q_b at time t_b , for an arbitrary Hamiltonian operator \underline{H} [which we write as $\underline{P}^2/2m + k \underline{H}_1(\underline{P}, \underline{Q}, t)$, \underline{H}_1

being arbitrary], can be written as a phase space path integral as follows:

$$K \equiv K_0 \int_{\mathcal{P}} dw(p, q) \exp \left\{ -\frac{ik}{\hbar} \int_T H_{no}[p(t), q(t), t] dt \right\} \quad (38)$$

$$\equiv K_0 \left\{ 1 + \sum_{j=1}^{\infty} \left(-\frac{ik}{\hbar} \right)^j \frac{1}{j!} \int_{\mathcal{P}} dw(p, q) \left[\int_T H_{no}(p(t), q(t), t) dt \right]^j \right\} \quad (39)$$

$$\equiv K_0 \left\{ 1 + \sum_{j=1}^{\infty} \left(-\frac{ik}{\hbar} \right)^j \frac{1}{j!} \int_{\mathcal{P}} dw(p, q) \int_T dt_1 H_{no}[p(t_1), q(t_1), t_1] \right. \\ \left. \dots \int_T dt_j H_{no}[p(t_j), q(t_j), t_j] \right\} \quad (40)$$

$$\equiv K_0 \left\{ 1 + \sum_{j=1}^{\infty} \left(-\frac{ik}{\hbar} \right)^j \frac{1}{j!} \int_T dt_1 \dots dt_j \right. \\ \times \lim_{(t_1 - t'_1) \rightarrow 0^+} \dots \lim_{(t_j - t'_j) \rightarrow 0^+} \int_{\mathcal{P}} dw(p, q) \\ \left. \times H_{no}[p(t'_1), q(t_1), t_1] \dots H_{no}[p(t'_j), q(t_j), t_j] \right\}, \quad (41)$$

where

(1) K_0 is the free-particle propagator (4), $w(p, q)$ is the free-particle measure defined by (1) and (2)ff, and $H_{no}(p, q, t)$ is a function obtained by

normal-ordering \widetilde{H}_1 (i.e. \widetilde{Q} before \widetilde{P}), then replacing \widetilde{Q} by q and \widetilde{P} by p ;

(2) The dot over the equal sign in the first three equations, (38)-(40), denotes a formal, as yet undefined, relation. The equations are undefined because as soon as one interchanges the path integral over \mathcal{P} and the time integral over T^J , one is faced with the path integral $\int_{\mathcal{P}} dw(p,q) H_{no}(p(t), q(t), t)$. Since the integrand contains terms coupling p and q at the same time t , this path integral is undefined because, as explained earlier, the pq correlation function, $\widetilde{G}(t, t')$ in (8), is undefined at $t = t'$. However, if we replace $\int_{\mathcal{P}} dw(p,q) H_{no}(p(t), q(t), t)$ by $\lim_{(t-t') \rightarrow 0^+} \int_{\mathcal{P}} dw(p,q) H_{no}(p(t'), q(t), t)$, the resulting expression is well-defined. This is done in the last equation, (41), which gives an unambiguous path-integral representation of K directly tied to the ordering of the factors in the quantum operator \widetilde{H}_1 ;

(3) It is not necessary that the H function in (38)-(41) be \widetilde{H}_{no} . However, if another is chosen, the "time" limits in (41) will be different. There is a close connection between the function chosen to replace \widetilde{H}_{no} and the type of time limits which will give a well-defined, correct expression for the propagator K . This will be proved and discussed in lemma 1 below;

(4) The propagator (41) satisfies the Schrödinger equation and the boundary condition:

$$\left[-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial q_b^2} + \widetilde{H}_1 \left(-i\hbar \frac{\partial}{\partial q_b}, q_b, t_b \right) - i\hbar \frac{\partial}{\partial t_b} \right] \langle q_b, t_b | q_a, t_a \rangle = 0 \quad (42a)$$

$$\langle q_b, t_b | q_a, t_a \rangle = \delta(q_b - q_a), \quad (42b)$$

to all orders in k ;

(5) The path integral in (41) can be evaluated in case a perturbation series in k is sought. The result, proved later in the paper, can be expressed in terms of only the classical Hamiltonian H_1 and the correspondence function F . It is:

$$K = K_0 \left[1 + \sum_{j=1}^{\infty} k^j \alpha_j \right], \quad (43)$$

where α_j is displayed in (70) and (71) below.

The proof of the theorem will consist in showing, by recurrence, that (41) satisfies (42) by using both (28) and (33) to relate the classical and quantum Hamiltonians. First, we give a simple illustration of the theorem.

Example

Calculate, to first order in k , the propagator corresponding to the following Hamiltonian:

$$\underline{H} = \underline{P}^2/2m + k \left[\alpha \underline{P} \underline{Q}^2 + \beta \underline{Q} \underline{P} \underline{Q} + (1-\alpha-\beta) \underline{Q}^2 \underline{P} \right], \quad (44)$$

(which represents all the possible \underline{H} s corresponding to $H_c = p^2/2m + kpq^2$).

Answer. The normal-ordered H is:

$$\underline{H} = \underline{P}^2/2m + k \left[\underline{Q}^2 \underline{P} - i\hbar(\beta + 2\alpha) \underline{Q} \right]. \quad (45)$$

Using (41), we have:

$$K = K_0 \left\{ 1 - \frac{i\hbar k}{T} \int_T dt \left[\lim_{t-t' \rightarrow 0^+} \int_{\mathcal{P}} q'(t) p(t') dw(p, q) - i\hbar(\beta + 2\alpha) \int_{\mathcal{P}} q(t) dw(p, q) \right] \right\} \quad (46)$$

$$= K_0 \left[1 - \frac{i\hbar k m}{3\hbar} (q_b^3 - q_a^3) + \frac{kT}{2} (q_b + q_a) (1 - \beta - 2\alpha) \right], \quad (47)$$

where we have used (25), (21), and (5)-(9). It can be directly verified that

(47) satisfies the Schrödinger equation to first order in k , i.e. that

$$\left\{ -(\hbar^2/2m) \frac{\partial^2}{\partial q_b^2} - i\hbar k \left[q_b^2 \frac{\partial}{\partial q_b} + (\beta - 2\alpha) q_b \right] - i\hbar \frac{\partial}{\partial t_b} \right\} K = O(k^2) \quad (48)$$

together with $\lim_{t_b \rightarrow t_a} K = \delta(q_b - q_a)$.

Note that we could have obtained the same result without normal-ordering first, as stated earlier. If we consider the function obtained by replacing \tilde{Q} and \tilde{P} by q and p in (44), order the times in the sequence suggested by (44), then take successive coincidence limits, we get:

$$K = K_0 \left\{ 1 - \frac{i\hbar k}{T} \int_T dt \left[\lim_{t-t' \rightarrow 0^+} \int_{\mathcal{P}} p(t) q^2(t') dw(p, q) + \beta \lim_{t-t' \rightarrow 0^+} \lim_{t'-t'' \rightarrow 0^+} \int_{\mathcal{P}} q(t) p(t') q(t'') dw(p, q) + (1 - \alpha - \beta) \lim_{t-t' \rightarrow 0^+} \int_{\mathcal{P}} q^2(t) p(t') dw(p, q) \right] \right\}, \quad (49)$$

which also yields the correct result (47). The general proof of this flexibility is found in lemma 1 below.

Proof of theorem

Lemma 1. For all phase-space functionals $F[q, p]$ which do not contain the path (p, q) evaluated at either t or t' , we have:

$$\lim_{t-t' \rightarrow 0^+} \int_{\mathcal{P}} F[q, p] [q(t)p(t') - q(t')p(t) - i\hbar] dw(p, q) = 0. \quad (50)$$

Proof. Consider the measure

$$dw_{tt'}(p, q) \equiv [q(t)p(t') - q(t')p(t) - i\hbar] dw(p, q). \quad (51)$$

Its Fourier transform is

$$\mathcal{F}w_{tt'}(\mu, \nu) = \int_{\mathcal{P}} e^{-i\langle \mu, q \rangle - i\langle \nu, p \rangle} dw_{tt'}(p, q). \quad (52)$$

To evaluate it, we proceed as follows:

$$\begin{aligned} & \int_{\mathcal{P}} q(t)p(t') e^{-i\langle \mu, q \rangle - i\langle \nu, p \rangle} dw(p, q) \\ &= -\frac{\partial^2}{\partial \lambda \partial \sigma} \int_{\mathcal{P}} e^{-i\langle \mu + \lambda \delta_t, q \rangle - i\langle \nu + \sigma \delta_{t'}, p \rangle} dw(p, q) \Big|_{\lambda=\sigma=0} \\ &= -\frac{\partial^2}{\partial \lambda \partial \sigma} \mathcal{F}w(\mu + \lambda \delta_t, \nu + \sigma \delta_{t'}) \Big|_{\lambda=\sigma=0} \end{aligned}$$

$$= F_w(\mu, \nu) \left[\frac{i\hbar}{2} G_{ab}(t, t') + \frac{i\hbar}{2} G_p(t, t') + i\hbar \bar{G}(t, t') + \xi(\mu, \nu; t, t') \right], \quad (53)$$

where

$$\begin{aligned} \xi(\mu, \nu; t, t') \equiv & \left[i\bar{q}(t) + \frac{i\hbar}{2} \int_T G_{ab}(t, s') d\mu(s') \right. \\ & \left. + \frac{i\hbar}{2} \int_T G_p(t, s') d\nu(s') + i\hbar \int_T \bar{G}(t, s') d\nu(s') \right] \\ & \times \left[i\bar{p}(t') + \frac{i\hbar}{2} \int_T G_{ab}(s, t') d\mu(s) + \frac{i\hbar}{2} \int_T G_p(s, t') d\nu(s) \right. \\ & \left. + i\hbar \int_T \bar{G}(s, t') d\mu(s) \right] \end{aligned} \quad (54)$$

and (2) was used. Therefore,

$$\begin{aligned} F_{w_{tt'}}(\mu, \nu) = F_w(\mu, \nu) \Big\{ & -i\hbar + \frac{i\hbar}{2} [G_{ab}(t, t') - G_{ab}(t', t)] \\ & + \frac{i\hbar}{2} [G_p(t, t') - G_p(t', t)] + i\hbar [\bar{G}(t, t') - \bar{G}(t', t)] \\ & + \xi(\mu, \nu; t, t') - \xi(\mu, \nu; t', t) \Big\}. \end{aligned} \quad (55)$$

Since G_{ab} , G_p , and ξ are continuous across the diagonal $t = t'$ (provided μ and ν are different from δ_t and $\delta_{t'}$), and since \bar{G} has a jump of magnitude 1 there [Eq. (27)], we conclude that

$$\lim_{t-t' \rightarrow 0^+} F_{w_{tt'}}(\mu, \nu) = 0. \quad (56)$$

Consequently, the measure $w_{tt'}(p,q)$ is effectively the zero measure in the limit $(t-t') \rightarrow 0^+$, provided (1) this limit is taken after a path integral with respect to $w_{tt'}(p,q)$ is performed [(50) is obviously false if the limit is taken before the path integral is done], and (2) the integrand does not contain p or q evaluated at t or t' [if it does, then $\xi(h,v;t,t')$ is no longer continuous across the diagonal $t = t'$]. Q.E.D.

This lemma insures that the various path integrals obtained by changing the form of the given Hamiltonian operator (by repeated use of the commutation relation $QP - PQ = i\hbar$) will all yield the same result. This was illustrated in the example above where the two path integrals (46) and (49), corresponding to the same Hamiltonian operator written in two different forms (45) and (44), gave the same correct answer.

Therefore, it is sufficient to prove the theorem for the normal-ordered form of H_1 , i.e. with H_{no} .

Lemma 2.

$$\begin{aligned} \left(\int_{t_a}^{t_b} f(t) dt \right)^n &= \int_{T^n} dt_1 \dots dt_n f(t_1) \dots f(t_n) \\ &= n! \int_{t_a}^{t_b} dt_1 \int_{t_a}^{t_1} dt_2 \dots \int_{t_a}^{t_{n-2}} dt_{n-1} \int_{t_a}^{t_{n-1}} dt_n f(t_1) \dots f(t_n). \end{aligned} \quad (57)$$

Proof. The lemma is true for $n = 1$. Assume it is true for $n = k - 1$.

Consider $F_k(s) \equiv \left(\int_{t_a}^s f(x) dx \right)^k$. Then

$$\dot{F}_k(s) = k F_{k-1}(s) f(s) = k f(s) (k-1)! \\ \times \int_{t_a}^s dt_1 \int_{t_a}^{t_1} dt_2 \dots \int_{t_a}^{t_{k-2}} dt_{k-1} f(t_1) \dots f(t_{k-1}). \quad (58)$$

Integrating from t_a to t_b with respect to s gives

$$F_k(t_b) = k! \int_{t_a}^{t_b} ds \int_{t_a}^s dt_1 \int_{t_a}^{t_1} dt_2 \dots \int_{t_a}^{t_{k-2}} dt_{k-1} f(t_1) \dots \\ f(t_{k-1}) f(s). \quad (59)$$

Changing variables: $s = t_1$, $t_1 = t_2$, ..., $t_{k-1} = t_k$, proves that the formula is true for $n = k$. Therefore the formula is proved true by recurrence. Q.E.D.

Proof to first order in k . The theorem will be proved by recurrence. Thus, we must first prove that (41) satisfies the Schrödinger equation (42) to first order in k . The propagator to first order in k is:

$$K = K_0 \left[1 - \frac{ik}{\hbar} \int_T dt \lim_{t-t' \rightarrow 0^+} \int_{\mathcal{P}} dw(p, q) H_{n_0}[p(t'), q(t), t] \right]. \quad (60)$$

Using the correspondence rule (28), along with (32) to put H_1 in normal-ordered form, we can write:

$$H_{n_0}[p(t'), q(t), t] = (2\pi\hbar)^{-2} \int dp dq du dv F(u, v, \hbar) H_1(p, q, t) \\ \times \exp \left\{ (i/\hbar) (qu + pv + uv/2) - iuq(t)/\hbar - ivp(t')/\hbar \right\}. \quad (61)$$

Substituting (61) in (60) reveals that the path integral is a particularly simple one, namely the Fourier transform of w at $(u\delta_t/\hbar, v\delta_t/\hbar)$. It can be evaluated using (2) and (5)-(9), and the result, after the limit, is:

$$K = K_0 [1 + \hbar \alpha_1], \quad (62)$$

where

$$\begin{aligned} \alpha_1 \equiv & -\frac{i}{\hbar(2\pi\hbar)^2} \int dp dq du dv F(u, v, \hbar) \int_T dt H_1(p, q, t) \\ & \times \exp \left\{ (i/\hbar)(qu + pv + uv/2) - (i/\hbar T) [uq_b(t-t_a) \right. \\ & + uq_a(t_b-t) + mv(q_b - q_a) + u^2(t-t_a)(t_b-t)/2m \\ & \left. + uv(t_b-t) - mv^2/2] \right\}. \end{aligned} \quad (63)$$

We must show that

$$\begin{aligned} & \left[-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial q_b^2} + \hbar \tilde{H}_1(q_b, -i\hbar \frac{\partial}{\partial q_b}, t_b) - i\hbar \frac{\partial}{\partial t_b} \right] \\ & \times [K_0(1 + \hbar \alpha_1)] = O(\hbar^2). \end{aligned} \quad (64)$$

Since K_0 satisfies the free-particle Schrödinger equation, we must simply show that the coefficient of \hbar is zero, i.e. that

$$\left[-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial q_b^2} - i\hbar \frac{\partial}{\partial t_b} \right] (K_0 \alpha_1) + \tilde{H}_1 K_0 = 0. \quad (65)$$

For \tilde{H}_1 we will use (33), its normal-ordered form in terms of the classical function H_1 . Thus,

$$\begin{aligned} \tilde{H}_1 K_0 = & (2\pi\hbar)^{-2} \int dp dq du dv F(u, v, \hbar) H_1(p, q, t_b) \\ & \times \left\{ \exp \left[\frac{i u}{\hbar} \left(q - q_b + \frac{v}{2} \right) + \frac{i p v}{\hbar} \right] \right\} \left(\frac{m}{2\pi i \hbar T} \right)^{1/2} \\ & \times \exp \left[\frac{i m}{2\hbar T} (q_b - v - q_a)^2 \right]. \end{aligned} \quad (66)$$

The remainder of the proof is tedious and straightforward. Differentiations with respect to q_b and t_b are performed under the integral sign, assuming interchangeability, using Leibnitz's rule, $(\partial/\partial t_b) \int_{t_a}^{t_b} \varphi(t, t_b) dt = \varphi(t_b, t_b) + \int_{t_a}^{t_b} [\partial \varphi(t, t_b) / \partial t_b] dt$, where needed. The $\tilde{H}_1 K_0$ term cancels the term equivalent to $\varphi(t_b, t_b)$. Upon collecting terms, the integrand vanishes, and the theorem is established to first order in \hbar for arbitrary Hamiltonian operators.

Proof to any order in \hbar . Using lemma 2 [Eq. (57)] along with (61), we can write K in (41) as:

$$K = K_0 \left[1 + \sum_{j=1}^{\infty} \hbar^j \alpha_j \right], \quad (67)$$

where

$$\begin{aligned}
\alpha_j \equiv & \left(-\frac{i}{\hbar}\right)^j \int_{t_a}^{t_b} dt_1 \int_{t_a}^{t_1} dt_2 \dots \int_{t_a}^{t_{j-2}} dt_{j-1} \int_{t_a}^{t_{j-1}} dt_j (2\pi\hbar)^{-2j} \int dq_1 \dots dq_j \\
& \times dp_1 \dots dp_j du_1 \dots du_j dv_1 \dots dv_j F(u_1, v_1, \hbar) \dots F(u_j, v_j, \hbar) \\
& \times H_1(p_1, q_1, t_1) \dots H_j(p_j, q_j, t_j) \\
& \times \exp \left[i \sum_{s=1}^j (q_s u_s + p_s v_s + u_s v_s / 2) / \hbar \right] \\
& \times \lim_{t_1 - t'_1 \rightarrow 0^+} \dots \lim_{t_j - t'_j \rightarrow 0^+} \int_{\mathcal{P}} dw(p, q) \exp \left\{ (-i/\hbar) [u_1 q(t_1) \right. \\
& \left. + \dots + u_j q(t_j) + v_1 p(t'_1) + \dots + v_j p(t'_j)] \right\}. \quad (68)
\end{aligned}$$

The path integral above is readily recognized as being

$$\mathcal{F}w \left[(-i/\hbar)(u_1 \delta_{t_1} + \dots + u_j \delta_{t_j}), (-i/\hbar)(v_1 \delta_{t'_1} + \dots + v_j \delta_{t'_j}) \right], \quad (69)$$

which can be evaluated, by use of (2) and (5)-(9). By virtue of lemma 2,

the times entering the integral are now ordered ($t_b \geq t_1 \geq t_2 \geq \dots \geq t_j \geq t_a$).

The result is:

$$\begin{aligned}
\alpha_j = & \left(-\frac{i}{\hbar}\right)^j \int_{t_a}^{t_b} dt_1 \int_{t_a}^{t_1} dt_2 \dots \int_{t_a}^{t_{j-2}} dt_{j-1} \int_{t_a}^{t_{j-1}} dt_j (2\pi\hbar)^{-2j} \\
& \times \int dq_1 \dots dq_j dp_1 \dots dp_j du_1 \dots du_j dv_1 \dots dv_j \\
& \times F(u_1, v_1, \hbar) \dots F(u_j, v_j, \hbar) H_1(p_1, q_1, t_1) \dots H_j(p_j, q_j, t_j) E_j, \quad (70)
\end{aligned}$$

where

$$\begin{aligned}
E_j \equiv \exp \Big\{ & (i/\hbar) \sum_{s=1}^j (q_s u_s + p_s v_s + u_s v_s / 2) \\
& - (i/\hbar T) \sum_{r=1}^j [u_r (t_r - t_a) q_b + u_r (t_b - t_r) q_a + m(q_b - q_a) v_r] \\
& - (i/\hbar T) \sum_{\substack{r,s=1 \\ r \geq s}}^j [u_r u_s (t_r - t_a)(t_b - t_s)/m - m v_r v_s / 2] \\
& - (i/\hbar T) \left[\sum_{\substack{r,s=1 \\ r \leq s}}^j u_r v_s (t_b - t_r) - \sum_{\substack{s=1, r=2 \\ r > s}}^j u_r v_s (t_r - t_a) \right] \Big\}.
\end{aligned} \tag{71}$$

We must now show that if

$$K_n \equiv K_0 \left[1 + \sum_{j=1}^n k^j \alpha_j \right] \tag{72}$$

satisfies the Schrödinger equation to n th order in k , then K_{n+1} satisfies it to $(n+1)$ th order in k , i.e.

$$\left(-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial q_b^2} + \hbar \tilde{H}_r - i\hbar \frac{\partial}{\partial t_b} \right) \left[K_0 \left(1 + \sum_{j=1}^{n+1} k^j \alpha_j \right) \right] = O(k^{n+2}). \tag{73}$$

This can be shown to be true if and only if the coefficient of k^{n+1} is 0, i.e. iff¹³

$$\left(-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial q_b^2} - i\hbar \frac{\partial}{\partial t_b} \right) (K_0 \alpha_{n+1}) + \underline{H}_1(K_0 \alpha_n) = 0 \quad (74)$$

(71) readily reduces to

$$\begin{aligned} \underline{H}_1(K_0 \alpha_n) - i\hbar K_0 \frac{\partial \alpha_{n+1}}{\partial t_b} - \frac{i\hbar K_0 (q_b - q_a)}{T} \frac{\partial \alpha_{n+1}}{\partial q_b} \\ - \frac{\hbar^2}{2m} K_0 \frac{\partial^2 \alpha_{n+1}}{\partial q_b^2} = 0. \end{aligned} \quad (75)$$

We will calculate each term in (75) separately and show that they cancel each other out. Using (33) to represent \underline{H}_1 , as we did earlier, we have:

$$\begin{aligned} \underline{H}_1(K_0 \alpha_n) &= (2\pi\hbar)^{-2} K_0 \int dp dq du dv F(u, v, \hbar) H_1(p, q, t_b) \\ &\times \exp \left\{ (i/\hbar) \left[uq + pv + uv/2 - uq_b - mv^2/2T - mv(q_b - q_a)/T \right] \right\} \\ &\times (-i/\hbar)^n \int_{t_a}^{t_b} dt_1 \int_{t_a}^{t_1} dt_2 \dots \int_{t_a}^{t_{n-1}} dt_n (2\pi\hbar)^{-2} \int dq_1 \dots dq_n dp_1 \dots dp_n \\ &\times du_1 \dots du_n dv_1 \dots dv_n F(u_1, v_1, \hbar) \dots F(u_n, v_n, \hbar) H_1(p_1, q_1, t_1) \\ &\times \dots H_1(p_n, q_n, t_n) \exp \left\{ (i/\hbar) \sum_{s=1}^n (q_s u_s + p_s v_s + u_s v_s / 2) \right. \\ &\quad - (i/\hbar T) \sum_{r=1}^n \left[u_r (t_r - t_a) (q_b - v) + u_r (t_b - t_r) q_a + m (q_b - v - q_a) v_r \right] \\ &\quad - (i/\hbar T) \sum_{\substack{r,s=1 \\ r \geq s}}^n \left[u_r u_s (t_r - t_a) (t_b - t_s) / m - m v_r v_s / 2 \right] \\ &\quad \left. - (i/\hbar T) \left[\sum_{\substack{r,s=1 \\ r \leq s}}^n u_r v_s (t_b - t_r) - \sum_{\substack{s=1, r=2 \\ r > s}}^n u_r v_s (t_r - t_a) \right] \right\}. \end{aligned} \quad (76)$$

The time-derivative term in (75), $-i\hbar K_0 \partial \alpha_{n+1} / \partial t_b$, can be written as A + B, where B is the derivative of the integrand and A evaluates the integrand at t_b . Thus,

$$\begin{aligned}
 A = & -i\hbar K_0 (-i/\hbar)^{n+1} \int_{t_a}^{t_b} dt_2 \cdots \int_{t_a}^{t_n} dt_{n+1} (2\pi\hbar)^{-2(n+1)} \\
 & \times \int dq_1 \cdots dq_{n+1} dp_1 \cdots dp_{n+1} du_1 \cdots du_{n+1} dv_1 \cdots dv_{n+1} \\
 & \times F(u_1, v_1, \hbar) \cdots F(u_{n+1}, v_{n+1}, \hbar) \\
 & \times H_1(p_1, q_1, t_b) H_1(p_2, q_2, t_2) \cdots H_1(p_{n+1}, q_{n+1}, t_{n+1}) \\
 & \times \exp \left\{ (i/\hbar) \sum_{s=1}^{n+1} (q_s u_s + p_s v_s + u_s v_s / 2) \right. \\
 & - (i/\hbar T) [u_1 T q_b + m(q_b - q_a) v_1 + \sum_{r=2}^{n+1} [u_r (t_r - t_a) q_b + u_r (t_b - t_r) q_a \\
 & \quad \left. + m(q_b - q_a) v_r]] \right. \\
 & - (i/\hbar T) \left[-m v_1^2 / 2 - m v_1 \sum_{r=2}^{n+1} v_r + \sum_{\substack{r,s=2 \\ r \geq s}}^{n+1} [u_r u_s (t_r - t_a)(t_b - t_s) / m \right. \\
 & \quad \left. - m v_r v_s / 2] \right] \\
 & \left. - (i/\hbar T) \left[\sum_{\substack{r,s=2 \\ r \leq s}}^{n+1} u_r v_s (t_b - t_r) - v_1 \sum_{r=2}^{n+1} u_r (t_r - t_a) \right. \right. \\
 & \quad \left. \left. - \sum_{\substack{s=2, r=3 \\ r > s}}^{n+1} u_r v_s (t_r - t_a) \right] \right\}. \quad (77)
 \end{aligned}$$

The t_b dependence of the integrand of $K_0 \alpha_{n+1}$ is of the form $\exp[at_b/(t_b - t_a)]$. Therefore,

$$\begin{aligned}
 B = & -i\hbar K_0 (-i/\hbar)^{n+1} \int_{t_a}^{t_b} dt_1 \dots \int_{t_a}^{t_b} dt_{n+1} (2\pi\hbar)^{2(n+1)} \\
 & \times \int dq_1 \dots dq_{n+1} dp_1 \dots dp_{n+1} du_1 \dots du_{n+1} dv_1 \dots dv_{n+1} \\
 & \times F(u_1, v_1, \hbar) \dots F(u_{n+1}, v_{n+1}, \hbar) \\
 & \times H_1(p_1, t_1, t_1) \dots H_1(p_{n+1}, q_{n+1}, t_{n+1}) E_{n+1} \\
 & \times (-i/\hbar T^2) \left\{ T q_a \sum_{r=1}^{n+1} u_r + \sum_{\substack{r,s=1 \\ r \geq s}}^{n+1} u_r u_s (t_r - t_a) T/m \right. \\
 & + T \sum_{\substack{r,s=1 \\ r \leq s}}^{n+1} u_r v_s - \sum_{r=1}^{n+1} [u_r (t_r - t_a) q_b + u_r (t_b - t_r) q_a \\
 & + m(q_b - q_a) v_r] - \sum_{\substack{r,s=1 \\ r \geq s}}^{n+1} [u_r u_s (t_r - t_a)(t_b - t_s)/m - m v_r v_s/2] \\
 & \left. - \sum_{\substack{r,s=1 \\ r \leq s}}^{n+1} u_r v_s (t_b - t_r) + \sum_{\substack{s=1, r=2 \\ r > s}}^{n+1} u_r v_s (t_r - t_a) \right\}. \quad (78)
 \end{aligned}$$

The q_b dependence of the integrand of α_{n+1} is of the simple form $\exp(aq_b)$. Therefore,

$$\begin{aligned}
 & -\frac{i\hbar}{T} K_0 (q_b - q_a) \frac{\partial \alpha_{n+1}}{\partial q_b} - \frac{\hbar^2}{2m} K_0 \frac{\partial^2 \alpha_{n+1}}{\partial q_b^2} \\
 & = i\hbar K_0 (-i/\hbar)^{n+1} \int_{t_a}^{t_b} dt_1 \cdots \int_{t_a}^{t_n} dt_{n+1} (2\pi\hbar)^{-2(n+1)} \\
 & \times \int dq_1 \cdots dq_{n+1} dp_1 \cdots dp_{n+1} du_1 \cdots du_{n+1} dv_1 \cdots dv_{n+1} \\
 & \times F(u_1, v_1, \hbar) \cdots F(u_{n+1}, v_{n+1}, \hbar) \\
 & \times H_1(p_1, q_1, t_1) \cdots H_1(p_{n+1}, q_{n+1}, t_{n+1}) E_{n+1} \\
 & \times (-i/\hbar T^2) \left\{ -(q_b - q_a) \sum_{r=1}^{n+1} [u_r(t_r - t_a) + m v_r] \right. \\
 & \quad \left. + \frac{1}{2m} \left[\sum_{r=1}^{n+1} (u_r(t_r - t_a) + m v_r) \right]^2 \right\}, \quad (79)
 \end{aligned}$$

where E_{n+1} is defined in (71).

From (76) and (77) one can show that $A + H_1(K_0 \alpha_n) = 0$. For this, it is sufficient to make the following changes of variable: in (77), $t_i = t_{i-1}$ for $i = 2$ to $n+1$; $C_i = C_{n+1}$ and $C_i = C_{i-1}$ for $i = 2$ to $n+1$, where c denotes u, v, p , or q ; in (76), $C = C_{n+1}$, where c denotes u, v, p , or q . It is also seen, by rearranging the terms inside the curly brackets of (78), that B added to either side of (79) gives zero.

The boundary condition (42b) is satisfied, as can be seen from the expression (43) for K . Indeed, K_0 satisfies (42b) and $\lim_{t_b \rightarrow t_a} \alpha_j = 0$, as can be seen from (71).

This completes the proof of the theorem. Q.E.D.

V. CONCLUSION

The object of this paper was to show that any Hamiltonian operator is amenable to a path-integral treatment, by providing an unambiguous, computationally viable formalism and taking proper account of the correspondence rule leading from the classical function to the quantum operator. The manner in which the correspondence rule is taken into account in a semiclassical expansion of the propagator (in powers of \hbar)¹⁴ will be the subject of a follow-up paper.

FOOTNOTES

1. Maurice M. Mizrahi, "The Weyl Correspondence and Path Integrals", J. Math. Phys. 16(1975), 2201-6.
2. L. Cohen, "Correspondence Rules and Path Integrals", J. Math. Phys. 17(1976), 597-8.
3. J. S. Dowker, "Path Integrals and Ordering Rules", J. Math. Phys. 17(1976), 1873-4.
4. Maurice M. Mizrahi, "Phase Space Path Integrals, Without Limiting Procedure", J. Math. Phys. 19(1978), 298-307.
5. L. Cohen, J. Math. Phys. 7 (1966), 781- .
6. However, if one is interested only in a perturbation series in α of the propagator corresponding to $H = [g(t) \underline{P}^2/2m + f(t) \underline{Q}^2/2 + k(t)(\underline{PQ} + \underline{QP})/2 + \alpha H_1(\underline{P}, \underline{Q}, t)]$, then the method described here applies directly.
7. A similar formalism obtained by different methods can be found in C. DeWitt-Morette, A. Maheshwari, and B. Nelson, "Path Integration in Phase Space", Gen. Rel. and Grav. 8(1977), 581-93.

8. Maurice M. Mizrahi, "On Path Integral Solutions of the Schrödinger Equation, Without Limiting Procedure", J. Math. Phys. 17(1976), 566-75.

9. Maurice M. Mizrahi, "Generalized Hermite Polynomials", J. Comp. and Appl. Math. 1(1975), 273-7.

10. The proof of (19) is similar to the one given in Ref. 8.

11. This scheme for correspondence rules was also used in M. M. Mizrahi, "On the Semiclassical Expansion in Quantum Mechanics for Arbitrary Hamiltonians", J. Math. Phys. 18(1977), 786-90, to determine the range of validity of a commonly-used formula for the WKB approximation of the propagator $\langle q_b, t_b | q_a, t_a \rangle$.

12. This formula can be proved by left-multiplying both sides of (28) by $\exp [i(\underline{Q}u' + \underline{P}v')/\hbar]$, taking traces [Note: $\text{tr } A \equiv \int dq \langle q | A | q \rangle$], and inserting complete sets of states after using Eq. (32).

13. In the k -expansion of the left side of (73), the vanishing of the 0th order term results from the Schrödinger equation for K_0 , the vanishing of the coefficient of k results from (65), and the vanishing of the coefficients of k^2 to k^{n+1} is what (74) (from $n=2$ to $n+1$) proves.

14. As discussed in Ref. 4, section IVB, a semiclassical expansion of a propagator which allows such an expansion [see (Ref. 11)] yields:

$$K = K_{WKB} \int_{\mathcal{P}_0} dw(p_0, q_0) \exp \left\{ (i/\hbar) \Omega[p_0, q_0] \right\}$$

where K_{WKB} is the semiclassical approximation to K , \mathcal{P}_0 is the same as \mathcal{P} except that $q_a = q_b = 0$, $(p_0, q_0) \in \mathcal{P}_0$,

$$\Omega[p_0, q_0] \equiv - \sum_{n=3}^{\infty} \sum_{k=0}^n \frac{n!}{(n-k)!k!} \int_T dt \left[\frac{\partial^n H_c}{\partial p^k \partial q^{n-k}} \right]_{\substack{q=q_c \\ p=p_c}}(t) p_0^k(t) q_0^{n-k}(t),$$

and w absorbs the full quadratic part of the expansion of the action functional about the classical path (q_c, p_c) . However, if one expands the exponential and attempts to carry out the path integral before the time integral, the indefiniteness discussed in this paper appears again. One conjectured answer, which remains to be verified, is that the $p_0^k(t) q_0^{n-k}(t)$ term should be "time-ordered" (in the manner discussed in this paper) with the same correspondence rule as the original Hamiltonian operator of the problem.

CNA Professional Papers — 1973 to Present*

- PP 103
Friedheim, Robert L., "Political Aspects of Ocean Ecology," 48 pp., Feb 1973, published in *Who Protects the Oceans*, John Lawrence Hargrove (ed.) (St. Paul: West Publ'g. Co., 1974), published by the American Society of International Law AD 757 936
- PP 104
Schick, Jack M., "A Review of James Cable, Gunboat Diplomacy: Political Applications of Limited Naval Forces," 8 pp., Feb 1973, (Reviewed in the *American Political Science Review*, Vol. LXVI, Dec 1972)
- PP 105
Corn, Robert J. and Phillips, Gary R., "On Optimal Correction of Gunfire Errors," 22 pp., Mar 1973, AD 761 674
- PP 106
Stoloff, Peter H., "User's Guide for Generalized Factor Analysis Program (FACTAN)," 35 pp., Feb 1973, (Includes an addendum published Aug 1974) AD 758 824
- PP 107
Stoloff, Peter H., "Relating Factor Analytically Derived Measures to Exogenous Variables," 17 pp., Mar 1973, AD 758 820
- PP 108
McConnell, James M. and Kelly, Anne M., "Superpower Naval Diplomacy in the Indo-Pakistani Crisis," 14 pp., 5 Feb 1973, (Published, with revisions, in *Survival*, Nov/Dec 1973) AD 761 675
- PP 109
Berghofer, Fred G., "Salaries: A Framework for the Study of Trends," 8 pp., Dec 1973, (Published in *Review of Income and Wealth*, Series 18, No. 4, Dec 1972)
- PP 110
Augusta, Joseph, "A Critique of Cost Analysis," 9 pp., Jul 1973, AD 766 376
- PP 111
Herrick, Robert W., "The USSR's 'Blue Belt of Defense' Concept: A Unified Military Plan for Defense Against Seaborne Nuclear Attack by Strike Carriers and Polaris/Poseidon SSBNs," 18 pp., May 1973, AD 766 375
- PP 112
Ginsberg, Lawrence H., "ELF Atmosphere Noise Level Statistics for Project SANGUINE," 29 pp., Apr 1974, AD 786 969
- PP 113
Ginsberg, Lawrence H., "Propagation Anomalies During Project SANGUINE Experiments," 5 pp., Apr 1974, AD 786 968
- PP 114
Maloney, Arthur P., "Job Satisfaction and Job Turnover," 41 pp., Jul 1973, AD 768 410
- PP 115
Silverman, Lester P., "The Determinants of Emergency and Elective Admissions to Hospitals," 145 pp., 18 Jul 1973, AD 766 377
- PP 116
Rehm, Allan S., "An Assessment of Military Operations Research in the USSR," 19 pp., Sep 1973, (Reprinted from *Proceedings, 30th Military Operations Research Symposium* (U. Secret Dec 1972) AD 770 116
- PP 117
McWhite, Peter B. and Ratliff, H. Donald, "Defending a Logistics System Under Mining Attack," 24 pp., Aug 1976 (to be submitted for publication in *Naval Research Logistics Quarterly*), presented at 44th National Meeting, Operations Research Society of America, November 1973, AD A030 454
*University of Florida
**Research supported in part under Office of Naval Research Contract N00014-68-0273-0017
- PP 118
Barfoot, C. Bernard, "Markov Duels," 18 pp., Apr 1973, (Reprinted from *Operations Research*, Vol. 22, No. 2, Mar-Apr 1974)
- PP 119
Stoloff, Peter and Lockman, Robert F., "Development of Navy Human Relations Questionnaire," 2 pp., May 1974, (Published in *American Psychological Association Proceedings*, 81st Annual Convention, 1973) AD 779 240
- PP 120
Smith, Michael W. and Schrimper, Ronald A., "Economic Analysis of the Intracity Dispersion of Criminal Activity," 30 pp., Jun 1974, (Presented at the Econometric Society Meetings, 30 Dec 1973) AD 780 538
*Economics, North Carolina State University
- PP 121
Devine, Eugene J., "Procurement and Retention of Navy Physicians," 21 pp., Jun 1974, (Presented at the 49th Annual Conference, Western Economic Association, Las Vegas, Nev., 10 Jun 1974) AD 780 539
- PP 122
Kelly, Anne M., "The Soviet Naval Presence During the Iraq-Kuwait Border Dispute: March-April 1973," 34 pp., Jun 1974, (Published in *Soviet Naval Policy*, ed. Michael McGwire, New York: Praeger) AD 780 592
- PP 123
Petersen, Charles C., "The Soviet Port-Clearing Operation in Bangladesh, March 1972-December 1973," 35 pp., Jun 1974, (Published in Michael McGwire, et al. (eds) *Soviet Naval Policy: Objectives and Constraints*, (New York: Praeger Publishers, 1974) AD 780 540
- PP 124
Friedheim, Robert L. and Jehn, Mary E., "Anticipating Soviet Behavior at the Third U.N. Law of the Sea Conference: USSR Positions and Dilemmas," 37 pp., 10 Apr 1974, (Published in *Soviet Naval Policy*, ed. Michael McGwire, New York: Praeger) AD 783 701
- PP 125
Weinland, Robert G., "Soviet Naval Operations—Ten Years of Change," 17 pp., Aug 1974, (Published in *Soviet Naval Policy*, ed. Michael McGwire, New York: Praeger) AD 783 962
- PP 126 — Classified.
- PP 127
Dragnich, George S., "The Soviet Union's Quest for Access to Naval Facilities in Egypt Prior to the June War of 1967," 64 pp., Jul 1974, AD 786 318
- PP 128
Stoloff, Peter and Lockman, Robert F., "Evaluation of Naval Officer Performance," 11 pp., (Presented at the 82nd Annual Convention of the American Psychological Association, 1974) Aug 1974, AD 784 012
- PP 129
Holen, Arlene and Horowitz, Stanley, "Partial Unemployment Insurance Benefits and the Extent of Partial Unemployment," 4 pp., Aug 1974, (Published in the *Journal of Human Resources*, Vol. IX, No. 3, Summer 1974) AD 784 010
- PP 130
Dismukes, Bradford, "Roles and Missions of Soviet Naval General Purpose Forces in Wartime: Pro-SSBN Operation," 20 pp., Aug 1974, AD 786 320
- PP 131
Weinland, Robert G., "Analysis of Gorskov's *Navies in War and Peace*," 45 pp., Aug 1974, (Published in *Soviet Naval Policy*, ed. Michael McGwire, New York: Praeger) AD 786 319
- PP 132
Kleinman, Samuel D., "Racial Differences in Hours Worked in the Market: A Preliminary Report," 77 pp., Feb 1975, (Paper read on 26 Oct 1974 at Eastern Economic Association Convention in Albany, N.Y.) AD A 005 517
- PP 133
Squires, Michael L., "A Stochastic Model of Regime Change in Latin America," 42 pp., Feb 1975, AD A 007 912
- PP 134
Root, R. M. and Cuniff, P. F., "A Study of the Shock Spectrum of a Two-Degree-of-Freedom Non-linear Vibratory System," 39 pp., Dec 1975, (Published in the condensed version of *The Journal of the Acoustic Society*, Vol. 60, No. 6, Dec 1976, pp. 1314
*Department of Mechanical Engineering, University of Maryland
- PP 135
Goudreau, Kenneth A., Kuzmack, Richard A., Wiedemann, Karen, "Analysis of Closure Alternatives for Naval Stations and Naval Air Stations," 47 pp., 3 Jun 1975 (Reprinted from "Hearing before the Subcommittee on Military Construction of the Committee on Armed Services," U.S. Senate, 93rd Congress, 1st Session, Part 2, 22 Jun 1973)
- PP 136
Strallings, William, "Cybernetics and Behavior Therapy," 13 pp., Jun 1975
- PP 137
Petersen, Charles C., "The Soviet Union and the Reopening of the Suez Canal: Mineclearing Operations in the Gulf of Suez," 30 pp., Aug 1975, AD A 015 376

*CNA Professional Papers with an AD number may be obtained from the National Technical Information Service, U.S. Department of Commerce, Springfield, Virginia 22151. Other papers are available from the author at the Center for Naval Analyses, 1401 Wilson Boulevard, Arlington, Virginia 22209.

CNA Professional Papers — 1973 to Present (Continued)

- PP 138
Stallings, William, "BRIDGE: An Interactive Dialogue-Generation Facility," 5 pp., Aug 1975 (Reprinted from IEEE Transactions on Systems, Man, and Cybernetics, Vol. 5, No. 3, May 1975)
- PP 139
Morgan, William F., Jr., "Beyond Folklore and Fables in Forestry to Positive Economics," 14 pp., (Presented at Southern Economic Association Meetings November, 1974) Aug 1975, AD A 015 293
- PP 140
Mahoney, Robert and Druckman, Daniel*, "Simulation, Experimentation, and Context," 36 pp., 1 Sep 1975. (Published in Simulation & Games, Vol. 6, No. 3, Sep 1975)
*Mathematica, Inc.
- PP 141
Mizrahi, Maurice M., "Generalized Hermite Polynomials," 5 pp., Feb 1976 (Reprinted from the Journal of Computational and Applied Mathematics, Vol. 1, No. 4 (1975), 273-277)
*Research supported by the National Science Foundation
- PP 142
Lockman, Robert F., Jehn, Christopher, and Shugart, William F. II., "Models for Estimating Premature Losses and Recruiting District Performance," 36 pp., Dec 1975 (Presented at the RAND Conference on Defense Manpower, Feb 1976, to be published in the conference proceedings)
AD A 020 443
- PP 143
Horowitz, Stanley and Sherman, Allan (LCDR, USN), "Maintenance Personnel Effectiveness in the Navy," 33 pp., Jan 1976 (Presented at the RAND Conference on Defense Manpower, Feb 1976, to be published in the conference proceedings) AD A021 581
- PP 144
Durch, William J., "The Navy of the Republic of China - History, Problems, and Prospects," 66 pp., Aug 1976 (To be published in "A Guide to Asiatic Fleets," ed by Barry M. Blechman, Naval Institute Press) AD A030 460
- PP 145
Kelly, Anne M., "Port Visits and the "Internationalist Mission" of the Soviet Navy," 36 pp., Apr 1976
AD A023 436
- PP 146
Palmour, Vernon E., "Alternatives for Increasing Access to Scientific Journals," 6 pp., Apr 1975 (Presented at the 1975 IEEE Conference on Scientific Journals, Cherry Hill, N.C., Apr 28-30; published in IEEE Transactions on Professional Communication, Vol. PC-18, No. 3, Sep 1975) AD A021 798
- PP 147
Kessler, J. Christian, "Legal Issues in Protecting Offshore Structures," 33 pp., Jun 1976 (Prepared under task order N00014-68-A-0091-0023 for ONR)
AD A028 389
- PP 148
McConnell, James M., "Military Political Tasks of the Soviet Navy in War and Peace," 62 pp., Dec 1975 (Published in Soviet Oceans Development Study of Senate Commerce Committee October 1976)
AD A022 590
- PP 149
Squires, Michael L., "Counterforce Effectiveness: A Comparison of the Tsipis "K" Measure and a Computer Simulation," 24 pp., Mar 1976 (Presented at the International Study Association Meetings, 27 Feb 1976) AD A022 591
- PP 150
Kelly, Anne M. and Petersen, Charles, "Recent Changes in Soviet Naval Policy: Prospects for Arms Limitations in the Mediterranean and Indian Ocean," 28 pp., Apr 1976, AD A 023 723
- PP 151
Horowitz, Stanley A., "The Economic Consequences of Political Philosophy," 8 pp., Apr 1976 (Reprinted from Economic Inquiry, Vol. XIV, No. 1, Mar 1976)
- PP 152
Mizrahi, Maurice M., "On Path Integral Solutions of the Schrodinger Equation, Without Limiting Procedure," 10 pp., Apr 1976 (Reprinted from Journal of Mathematical Physics, Vol. 17, No. 4 (Apr 1976), 566-575)
*Research supported by the National Science Foundation
- PP 153
Mizrahi, Maurice M., "WKB Expansions by Path Integrals, With Applications to the Anharmonic Oscillator," 137 pp., May 1976 (Submitted for publication in Annals of Physics) AD A023 440
*Research supported by the National Science Foundation
- PP 154
Mizrahi, Maurice M., "On the Semi-Classical Expansion in Quantum Mechanics for Arbitrary Hamiltonians," 19 pp., May 1976 (To appear in the Journal of Mathematical Physics) AD A025 441
- PP 155
Squires, Michael L., "Soviet Foreign Policy and Third World Nations," 26 pp., Jun 1976 (Prepared for presentation at the Midwest Political Science Association meetings, Apr 30, 1976) AD A028 388
- PP 156
Stallings, William, "Approaches to Chinese Character Recognition," 12 pp., Jun 1976 (Reprinted from Pattern Recognition (Pergamon Press), Vol. 8, pp. 87-98, 1976) AD A028 692
- PP 157
Morgan, William F., "Unemployment and the Pentagon Budget: Is There Anything in the Empty Fuel Barrel?" 20 pp., Aug 1976 AD A030 455
- PP 158
Haskell, LCDR Richard D. (USN), "Experimental Validation of Probability Predictions," 28 pp., Aug 1976 (Presented at the Military Operations Research Society Meeting, Fall 1976) AD A030 458
- PP 159
McConnell, James M., "The Gorskoy Articles, The New Gorskoy Book and Their Relation to Policy," 93 pp., Jul 1976 (To be printed in Soviet Naval Influence: Domestic and Foreign Dimensions, ed by M. McGwire and J. McDonnell, New York: Praeger)
AD A029 227
- PP 160
Wilson, Desmond P., Jr., "The U.S. Sixth Fleet and the Conventional Defense of Europe," 50 pp., Sep 1976 (Submitted for publication in Adelphi Papers, I.I.S.S., London) AD A030 457
- PP 161
Melich, Michael E. and Peet, Vice Adm. Ray (USN, Retired), "Fleet Commanders: Afloat or Ashore?" 9 pp., Aug 1976 (Reprinted from U.S. Naval Institute Proceedings, Jun 1976) AD A030 456
- PP 162
Friedheim, Robert L., "Parliamentary Diplomacy," 106 pp. Sep 1976 AD A033 306
- PP 163
Lockman, Robert F., "A Model for Predicting Recruit Losses," 9 pp., Sep 1976 (Presented at the 84th annual convention of the American Psychological Association, Washington, D.C., 4 Sep 1976)
AD A030 459
- PP 164
Mahoney, Robert B., Jr., "An Assessment of Public and Elite Perceptions in France, The United Kingdom, and the Federal Republic of Germany," 31 pp., Feb 1977 (Presented at Conference "Perception of the U.S. - Soviet Balance and the Political Uses of Military Power" sponsored by Director, Advanced Research Projects Agency, April 1976) AD 036 509
- PP 165
Jondrow, James M., "Effects of Trade Restrictions on Imports of Steel," 67 pp., November 1976, (Delivered at ILAB Conference in Dec 1976)
- PP 166
Feldman, Paul, "Impediments to the Implementation of Desirable Changes in the Regulation of Urban Public Transportation," 12 pp., Oct 1976,
AD A033 322
- PP 166 - Revised
Feldman, Paul, "Why It's Difficult to Change Regulation," Oct 1976
- PP 167
Kleinman, Samuel, "ROTC Service Commitments: A Comment," 4 pp., Nov 1976, (To be published in Public Choice, Vol. XXIV, Fall 1976) AD A033 305
- PP 168
Lockman, Robert F., "Revalidation of CNA Support Personnel Selection Measures," 36 pp., Nov 1976
- PP 169
Jacobson, Louis S., "Earnings Losses of Workers Displaced from Manufacturing Industries," 38 pp., Nov 1976, (Delivered at ILAB Conference in Dec 1976)
- PP 170
Brechtling, Frank P., "A Time Series Analysis of Labor Turnover," Nov 1976, (Delivered at ILAB Conference in Dec 1976)
- PP 171
Ralston, James M., "A Diffusion Model for GaP Red LED Degradation," 10 pp., Nov 1976, (Published in Journal of Applied Physics, Vol. 47, pp. 4518-4522, Oct 1976)
- PP 172
Classen, Kathleen P., "Unemployment Insurance and the Length of Unemployment," Dec 1976, (Presented at the University of Rochester Labor Workshop on 16 Nov 1976)
- PP 173
Kleinman, Samuel D., "A Note on Racial Differences in the Added Worker/ Discouraged Worker Controversy," 2 pp., Dec 1976, (Published in the American Economist, Vol. XX, No. 1, Spring 1976)

CNA Professional Papers — 1973 to Present (Continued)

- PP 174
Mahoney, Robert B., Jr., "A Comparison of the Brookings and International Incidents Projects," 12 pp. Feb 1977 AD 037 206
- PP 175
Lestne, Daniel; Stoloff, Peter and Spruill, Nancy, "Public Drug Treatment and Addict Crime," June 1976, (Published in Journal of Legal Studies, Vol. 5, No. 2)
- PP 176
Felix, Wendt, "Correlates of Retention and Promotion for USNA Graduates," 38 pp., Mar 1977
- PP 177
Lockman, Robert F. and Warner, John T., "Predicting Attrition: A Test of Alternative Approaches," 33 pp. Mar 1977. (Presented at the OSD/ONR Conference on Enlisted Attrition Xerox International Training Center, Leesburg, Virginia, 4-7 April 1977)
- PP 178
Kleinman, Samuel D., "An Evaluation of Navy Unrestricted Line Officer Accession Programs," 23 pp. April 1977, (To be presented at the NATO Conference on Manpower Planning and Organization Design, Sirena, Italy, 20 June 1977)
- PP 179
Stoloff, Peter H. and Balut, Stephen J., "Vacate: A Model for Personnel Inventory Planning Under Changing Management Policy," 14 pp. April 1977, (To be presented at the NATO Conference on Manpower Planning and Organization Design, Sirena, Italy, 20 June 1977)
- PP 180
Horowitz, Stanley A. and Sherman, Allan, "The Characteristics of Naval Personnel and Personnel Performance," 16 pp. April 1977, (To be presented at the NATO Conference on Manpower Planning and Organization Design, Sirena, Italy, 20 June 1977)
- PP 181
Balut, Stephen J. and Stoloff, Peter, "An Inventory Planning Model for Navy Enlisted Personnel," 35 pp., May 1977, (Prepared for presentation at the Joint National Meeting of the Operations Research Society of America and The Institute for Management Science, 9 May 1977, San Francisco, California)
- PP 182
Murray, Russell, 2nd, "The Quest for the Perfect Study or My First 1138 Days at CNA," 57 pp., April 1977
- PP 183
Kassing, David, "Changes in Soviet Naval Forces," 33 pp., November, 1976, (To be published as a chapter in a book published by The National Strategic Information Center)
- PP 184
Lockman, Robert F., "An Overview of the OSD/ONR Conference on First Term Enlisted Attrition," 22 pp., June 1977, (Presented to the 39th MORS Working Group on Manpower and Personnel Planning, Annapolis, Md., 28-30 June 1977)
- PP 185
Kassing, David, "New Technology and Naval Forces in the South Atlantic," 22 pp. (This paper was the basis for a presentation made at the Institute for Foreign Policy Analyses, Cambridge, Mass., 28 April 1977.)
- PP 186
Mizrahi, Maurice M., "Phase Space Integrals, Without Limiting Procedure," 31 pp., May 1977, (Submitted for publication in Journal of Mathematical Physics)
- PP 187
Code, Russell C., "Nomography for Operations Research," 35 pp., April 1977, (Presented at the Joint National Meeting of the Operations Research Society of America and The Institute for Management Services, San Francisco, California, 9 May 1977)
- PP 188
Dorch, William J., "Information Processing and Outcome Forecasting for Multilateral Negotiations: Testing One Approach," 53 pp., May 1977 (Prepared for presentation to the 18th Annual Convention of the International Studies Association, Chase Park Plaza Hotel, St. Louis, Missouri, March 16-20, 1977)
- PP 189
Code, Russell C., "Error Detection in Computerized Information Retrieval Data Bases," July, 1977, 13 pp. Presented at the Sixth Cranfield International Conference on Mechanized Information Storage and Retrieval Systems, Cranfield Institute of Technology, Cranfield, Bedford, England, 26-29 July 1977
- PP 190
Mahoney, Robert B., Jr., "European Perceptions and East-West Competition," 96 pp., July 1977 (Prepared for presentation at the annual meeting of the International Studies Association, St. Louis, Mo., March, 1977)
- PP 191
Sawyer, Ronald, "The Independent Field Assignment: One Man's View," August 1977, 25 pp.
- PP 192
Holen, Arlene, "Effects of Unemployment Insurance Entitlement on Duration and Job Search Outcome," August 1977, 6 pp., (Reprinted from Industrial and Labor Relations Review, Vol. 30, No. 4, Jul 1977)
- PP 193
Horowitz, Stanley A., "A Model of Unemployment Insurance and the Work Test," August 1977, 7 pp. (Reprinted from Industrial and Labor Relations Review, Vol. 30, No. 40, Jul 1977)
- PP 194
Classen, Kathleen P., "The Effects of Unemployment Insurance on the Duration of Unemployment and Subsequent Earnings," August 1977, 7 pp. (Reprinted from Industrial and Labor Relations Review, Vol. 30, No. 40, Jul 1977)
- PP 195
Brechtling, Frank, "Unemployment Insurance Taxes and Labor Turnover: Summary of Theoretical Findings," 12 pp. (Reprinted from Industrial and Labor Relations Review, Vol. 30, No. 40, Jul 1977)
- PP 196
Ralston, J. M. and Lorimer, O. G., "Degradation of Bulk Electroluminescent Efficiency in Zn, O-Doped GaP LED's," July 1977, 3 pp. (Reprinted from IEEE Transactions on Electron Devices, Vol. ED-24, No. 7, July 1977)
- PP 197
Wells, Anthony R., "The Centre for Naval Analyses," 14 pp., Dec 1977
- PP 198
Classen, Kathleen P., "The Distributional Effects of Unemployment Insurance," 25 pp., Sept. 1977 (Presented at a Hoover Institution Conference on Income Distribution, Oct 7-8, 1977)
- PP 199
Dorch, William J., "Revolution From A F.A.R. — The Cuban Armed Forces in Africa and the Middle East," Sep 1977, 16 pp.
- PP 200
Powers, Bruce F., "The United States Navy," 40 pp. Dec 1977. (To be published in American Military Machine)
- PP 201
Dorch, William J., "The Cuban Military in Africa and the Middle East: From Algeria to Angola," Sep 1977, 67 pp.
- PP 202
Feldman, Paul, "Why Regulation Doesn't Work," (Reprinted from *Technological Change and Welfare in the Regulated Industries and Review of Social Economy*, Vol. XXIX, March, 1971, No. 1.) Sep 1977, 8 pp.
- PP 203
Feldman, Paul, "Efficiency, Distribution, and the Role of Government in a Market Economy," (Reprinted from *The Journal of Political Economy*, Vol. 79, No. 3, May/June 1971.) Sep 1977, 19 pp.
- PP 204
Wells, Anthony R., "The 1967 June War: Soviet Naval Diplomacy and The Sixth Fleet — A Reappraisal," Oct 1977, 36 pp.
- PP 205
Code, Russell C., "A Bibliometric Examination of the Square Root Theory of Scientific Publication Productivity," (Presented at the annual meeting of the American Society for Information Science, Chicago, Illinois, 29 September 1977.) Oct 1977, 6 pp.
- PP 206
McConnell, James M., "Strategy and Missions of the Soviet Navy in the Year 2000," 48 pp., Nov 1977, (To be presented at a Conference on Problems of Sea Power as we Approach the 21st Century, sponsored by the American Enterprise Institute for Public Policy Research, 6 October 1977, and subsequently published in a collection of papers by the Institute)

CNA Professional Papers — 1973 to Present (Continued)

PP 207

Goldberg, Lawrence, "Cost-Effectiveness of Potential Federal Policies Affecting Research & Development Expenditures in the Auto, Steel and Food Industries," 36 pp., Oct 1977, (Presented at Southern Economic Association Meetings beginning 2 November 1977)

PP 208

Roberts, Stephen S., "The Decline of the Overseas Station Fleets: The United States Asiatic Fleet and the Shanghai Crisis, 1932," 18 pp., Nov 1977, (Reprinted from The American Neptune, Vol. XXXVII, No. 3, July 1977)

PP 209 - Classified.

PP 210

Kassing, David, "Protecting The Fleet," 40 pp., Dec 1977 (Prepared for the American Enterprise Institute Conference on Problems of Sea Power as We Approach the 21st Century, October 6-7, 1977)

PP 211

Mizrahi, Maurice M., "On Approximating the Circular Coverage Function," 14 pp., Feb 1978

PP 213

Mangel, Marc, "Fluctuations in Systems with Multiple Steady States. Application to Lanchester Equations," 12 pp., Feb 78, (Presented at the First Annual Workshop on the Information Linkage Between Applied Mathematics and Industry, Naval PG School, Feb 23-25, 1978)

PP 214

Weinland, Robert G., "A Somewhat Different View of The Optimal Naval Posture," 37 pp., Jun 1978 (Presented at the 1976 Convention of the American Political Science Association (APSA)/US Panel on "Changing Strategic Requirements and Military Posture", Chicago, Ill., September 2, 1976)

PP 215

Coile, Russell C., "Comments on: Principles of Information Retrieval by Manfred Kochen, 10 pp., Mar 78, (Published as a Letter to the Editor, Journal of Documentation, Vol. 31, No. 4, pages 298-301, December 1975)

PP 216

Coile, Russell C., "Lotka's Frequency Distribution of Scientific Productivity," 18 pp., Feb 1978, (Published in the Journal of the American Society for Information Science, Vol. 28, NO. 6, pp. 366-370, November 1977)

PP 217

Coile, Russell C., "Bibliometric Studies of Scientific Productivity," 17 pp., Mar 78, (Presented at the Annual meeting of the American Society for Information Science held in San Francisco, California, October 1976.)

PP 218 - Classified.

PP 219

Huntzinger, R. LaVar, "Market Analysis with Rational Expectations: Theory and Estimation," 60 pp., Apr 78 (To be submitted for publication in Journal of Econometrics)

PP 220

Maurer, Donald E., "Diagonalization by Group Matrices," 26 pp., Apr 78

PP 221

Weinland, Robert G., "Superpower Naval Diplomacy in the October 1973 Arab-Israeli War," 76 pp., Jun 1978

PP 222

Mizrahi, Maurice M., "Correspondence Rules and Path Integrals," 30 pp., Jun 1978 (Submitted for Publication in The Journal of Mathematical Physics)